

THE h -VECTOR OF A LADDER DETERMINANTAL RING COGENERATED BY 2×2 MINORS IS LOG-CONCAVE

MARTIN RUBEY

ABSTRACT. We show that the h -vector of a ladder determinantal ring cogenerated by $M = [u_1 \mid v_1]$ is log-concave. Thus we prove an instance of a conjecture of Stanley, resp. Conca and Herzog.

In honour of Miriam Rubey, at the occasion of her second birthday

1. INTRODUCTION

Definition 1.1. A sequence of real numbers $a_1, a_2, \dots, a_n$ is *logarithmically concave*, for short *log-concave*, if $a_{i-1}a_{i+1} \leq a_i^2$ for $i \in \{2, 3, \dots, n-1\}$.

Numerous sequences arising in combinatorics and algebra have, or seem to have this property. In the paper [13] written in 1989, Richard Stanley collected various results on this topic. (For an update see [3].) There he also stated the following conjecture:

Conjecture 1.2. Let $R = R_0 \oplus R_1 \oplus \dots$ be a graded (Noetherian) Cohen-Macaulay (or perhaps Gorenstein) domain over a field $K = R_0$, which is generated by R_1 and has Krull dimension d . Let $H(R, m) = \dim_K R_m$ be the Hilbert function of R and write

$$\sum_{m \geq 0} H(R, m)x^m = (1-x)^{-d} \sum_{i=0}^s h_i x^i.$$

Then the sequence $h_0, h_1, \dots, h_s$ is log-concave.

The sequence $h_0, h_1, \dots, h_s$ is called the h -vector of the ring. Originally the question was to decide whether a given sequence can arise as the h -vector of some ring. In this sense the validity of the conjecture would imply that log-concavity was a necessary condition on the h -vector.

It is now known however [12, 3] that Stanley's conjecture is not true in general. Several natural weakenings have been considered, but are still open. For example, Aldo Conca and Jürgen Herzog conjectured that the h -vector would be log-concave for the special case where R is a ladder determinantal ring. (Note that ladder determinantal rings are Cohen-Macaulay, as was shown in [8, Corollary 4.10], but not necessarily Gorenstein.) We will prove the conjecture of Conca and Herzog in the simplest case, i.e., where R is a ladder determinantal ring cogenerated by 2×2 minors, see Corollary 4.6.

In the case of ladder determinantal rings the h -vector has a nice combinatorial interpretation. This follows from work of Abhyankar and Kulkarni [1, 2, 10, 11], Bruns, Conca, Herzog, and Trung [4, 5, 6, 8]. In the following paragraphs, which are taken almost verbatim from [9], we will explain these matters.

2. LADDERS, LADDER DETERMINANTAL RINGS AND NON-INTERSECTING LATTICE PATHS

First we have to introduce the notion of a ladder:

Definition 2.1. Let $\mathbf{X} = (x_{i,j})_{0 \leq i \leq b, 0 \leq j \leq a}$ be a $(b+1) \times (a+1)$ matrix of indeterminates. Let $\mathbf{Y} = (y_{i,j})_{0 \leq i \leq b, 0 \leq j \leq a}$ be another matrix of the same dimensions, with the property that $y_{i,j} \in \{0, x_{i,j}\}$, and if $y_{i,j} = x_{i,j}$ and $y_{i',j'} = x_{i',j'}$, where $i \leq i'$ and $j \leq j'$ then $y_{r,s} = x_{r,s}$ for all r and s with $i \leq r \leq i'$ and $j \leq s \leq j'$. Such a matrix $\mathbf{Y}$ is called a *ladder*.

A *ladder region* L is a subset of $\mathbb{Z}^2$ with the property that if (i, j) and $(i', j') \in L$, $i \leq i'$ and $j \geq j'$ then $(r, s) \in L$ for all $r \in \{i, i+1, \dots, i'\}$ and $s \in \{j', j'+1, \dots, j\}$. Clearly, a ladder region can be described by two weakly increasing functions $\underline{L}$ and $\overline{L}$, such that L is exactly the set of points $\{(i, j) : \underline{L}(i) \leq j \leq \overline{L}(i)\}$.

We associate with $\mathbf{Y}$ a ladder region $L \subset \mathbb{Z}^2$ via $(j, b-i) \in L$ if and only if $y_{i,j} = x_{i,j}$.

In Figure 1.a an example of a ladder with $a = 8$ and $b = 9$ is shown, the corresponding ladder region is shown in Figure 1.b.

Now we can define the ring we are dealing with:

Definition 2.2. Given a $(b+1) \times (a+1)$ matrix $\mathbf{Y}$ which is a ladder, fix a “bivector” $M = [u_1, u_2, \dots, u_n \mid v_1, v_2, \dots, v_n]$ of integers with $1 \leq u_1 < u_2 < \dots < u_n \leq b+1$ and $1 \leq v_1 < v_2 < \dots < v_n \leq a+1$. By convention we set $u_{n+1} = b+2$ and $v_{n+1} = a+2$.

Let $K[\mathbf{Y}]$ denote the ring of all polynomials over some field K in the $y_{i,j}$ ’s, where $0 \leq i \leq b$ and $0 \leq j \leq a$. Furthermore, let $I_M(\mathbf{Y})$ be the ideal in $K[\mathbf{Y}]$ that is generated by those $t \times t$ minors of $\mathbf{Y}$ that contain only nonzero entries, whose rows form a subset of the last $u_t - 1$ rows or whose columns form a subset of the last $v_t - 1$ columns, $t \in \{1, 2, \dots, n+1\}$. Thus, for $t = n+1$ the rows and columns of minors are unrestricted.

The ideal $I_M(\mathbf{Y})$ is called a *ladder determinantal ideal generated by the minors defined by M* . We call $R_M(\mathbf{Y}) = K[\mathbf{Y}]/I_M(\mathbf{Y})$ the *ladder determinantal ring cogenerated by the minors defined by M* , or, in abuse of language, the *ladder determinantal ring cogenerated by M* .

Note that we could restrict ourselves to the case $u_1 = v_1 = 1$, because all the elements of $\mathbf{Y}$ that are in one of the last $u_1 - 1$ rows or in one of the last $v_1 - 1$ columns are in the ideal.

Next, we introduce the combinatorial objects that will accompany us throughout the rest of this paper:

Definition 2.3. A *two-rowed array of length k* is a pair of strictly increasing sequences of integers, both of length k . A two-rowed array $T = \begin{pmatrix} a_1 & a_2 & \dots & a_k \\ b_1 & b_2 & \dots & b_k \end{pmatrix}$ is *bounded* by $A = (A_1, A_2)$ and $E = (E_1, E_2)$, if

$$\begin{aligned} A_1 &\leq a_1 < a_2 < \dots < a_k \leq E_1 - 1 \\ \text{and } A_2 + 1 &\leq b_1 < b_2 < \dots < b_k \leq E_2. \end{aligned}$$

Given *any* subset L of $\mathbb{Z}^2$, we say that the two-rowed array T is *in L* , if $(a_i, b_i) \in L$ for $i \in \{1, 2, \dots, k\}$. By $\mathcal{T}_k^L(A \mapsto E)$ we will denote the set of two-rowed arrays of length k , bounded by A and E which are in L . The *total length* of a family of two-rowed arrays is just the sum of the lengths of its members.

Let $T_1 = \begin{pmatrix} a_1 & a_2 & \dots & a_k \\ b_1 & b_2 & \dots & b_k \end{pmatrix}$ and $T_2 = \begin{pmatrix} x_1 & x_2 & \dots & x_l \\ y_1 & y_2 & \dots & y_l \end{pmatrix}$ be two-rowed arrays bounded by $A^{(1)} = (A_1^{(1)}, A_2^{(1)})$ and $E^{(1)} = (E_1^{(1)}, E_2^{(1)})$ and $A^{(2)} = (A_1^{(2)}, A_2^{(2)})$ and $E^{(2)} = (E_1^{(2)}, E_2^{(2)})$ respectively. Set $a_{k+1} = E_1^{(1)}$ and $b_0 = A_2^{(1)}$. We say that T_1 and T_2 *intersect* if there are indices I and J such that

$$(\times) \quad \begin{aligned} x_J &\leq a_I \\ b_{I-1} &\leq y_J \end{aligned}$$


where $1 \leq I \leq k+1$ and $1 \leq J \leq l$. A family of two-rowed arrays is *non-intersecting* if no two arrays in it intersect.

Note that a two-rowed array in $\mathcal{T}_k^L(A \mapsto E)$ can be visualized by a lattice path with east and north steps, that starts in A and terminates in E and has exactly k north-east turns which are all in L : Each pair (a_i, b_i) of a two-rowed array $\begin{pmatrix} a_1 & a_2 & \dots & a_k \\ b_1 & b_2 & \dots & b_k \end{pmatrix}$ then corresponds to a north-east turn of the lattice path. It is easy to see that Condition $(\times)$ holds if and only if the lattice paths corresponding to T_1 and T_2 intersect.

For an example see Figure 1.c, where the three two-rowed arrays

$$T^{(1)} = \begin{pmatrix} 2 & 3 \\ 6 & 7 \end{pmatrix}, T^{(2)} = \begin{pmatrix} 3 & 5 \\ 4 & 6 \end{pmatrix} \text{ and } T^{(3)} = \begin{pmatrix} 2 & 4 & 6 \\ 1 & 3 & 4 \end{pmatrix}$$

bounded by $A^{(1)} = (0, 3)$, $A^{(2)} = (0, 2)$, $A^{(3)} = (0, 0)$ and $E^{(1)} = (5, 9)$, $E^{(2)} = (7, 9)$, $E^{(3)} = (8, 9)$ are shown as lattice paths. The points of the ladder-region L are drawn as small dots, the circles indicate the start- and endpoints and the big dots indicate the north-east turns.



a. a ladder with $a = 8$
and $b = 9$

b. the corresponding
ladder region

c. a triple of non-
intersecting lattice
paths in this ladder

FIGURE 1.

3. A COMBINATORIAL INTERPRETATION OF THE h -VECTOR OF A LADDER DETERMINANTAL RING

We are now ready to state the theorem which reveals the combinatorial nature of the h -vector of $R_M(\mathbf{Y}) = K[\mathbf{Y}]/I_M(\mathbf{Y})$, the ladder determinantal ring cogenerated by M .

Theorem 3.1. *Let $\mathbf{Y} = (y_{i,j})_{0 \leq i \leq b, 0 \leq j \leq a}$ be a ladder and let $M = [u_1, u_2, \dots, u_n \mid v_1, v_2, \dots, v_n]$ be a bivector of integers with $1 \leq u_1 < u_2 < \dots < u_n \leq a+1$ and $1 \leq v_1 < v_2 < \dots < v_n \leq b+1$. For $i \in \{1, 2, \dots, n\}$ let*

$$\begin{aligned} A^{(i)} &= (0, u_{n+1-i} - 1) \\ E^{(i)} &= (a - v_{n+1-i} + 1, b). \end{aligned}$$

Let $L^{(n)} = L$ be the ladder region associated with $\mathbf{Y}$ and for $i \in \{1, 2, \dots, n-1\}$ let

$$L^{(i)} = \{(x, y) \in L^{(i+1)} : x \leq E_1^{(i)}, y \geq A_2^{(i)} \text{ and } (x+1, y-1) \in L^{(i+1)}\}.$$

Finally, for $i \in \{1, 2, \dots, n\}$ let

$$B^{(i)} = \{(x, y) \in L^{(i)} : (x+1, y-1) \notin L^{(i)}\}.$$

and let d be the cardinality of $\bigcup_{i=1}^n B^{(i)}$.

Then, under the assumption that all of the points $A^{(i)}$ and $E^{(i)}$, $i \in \{1, 2, \dots, n\}$, lie inside the ladder region L , the Hilbert series of the ladder determinantal ring $R_M(\mathbf{Y}) = K[\mathbf{Y}]/I_M(\mathbf{Y})$ equals

$$\sum_{\ell \geq 0} \dim_K R_M(\mathbf{Y})_\ell z^\ell = \frac{\sum_{\ell \geq 0} |\mathcal{T}_\ell^L(\mathbf{A} \mapsto \mathbf{E})| z^\ell}{(1-z)^d}.$$

Here, $R_M(\mathbf{Y})_\ell$ denotes the homogeneous component of degree ℓ in $R_M(\mathbf{Y})$ and $|\mathcal{T}_\ell^L(\mathbf{A} \mapsto \mathbf{E})|$ is the number of non-intersecting families of two-rowed arrays with total length ℓ , such that the i^{th} two-rowed array is bounded by $A^{(i)}$ and $E^{(i)}$ and is in $L^{(i)} \setminus B^{(i)}$ for $i \in \{1, 2, \dots, n\}$.


The sets $B^{(i)}$, $i \in \{1, 2, \dots, n\}$ can be visualized as being the lower-right boundary of $L^{(i)}$. Viewed as a path, there are exactly $E_1^{(i)} - A_1^{(i)} + E_2^{(i)} - A_2^{(i)} + 1$ lattice points on $B^{(i)}$, but not all of them are necessarily in L . However, if L is an upper ladder, that is, $(a, 0) \in L$, then this must be the case and we have

$$\begin{aligned} d &= \sum_{i=1}^n \left(E_1^{(i)} - A_1^{(i)} + E_2^{(i)} - A_2^{(i)} + 1 \right) \\ &= \sum_{i=1}^n (a - v_{n+1-i} + 1 + b - u_{n+1-i} + 1 + 1) \\ &= n(a + b + 3) - \sum_{i=1}^n (u_i + v_i), \end{aligned}$$

as in [9].

In Figure 2.a, an example for a ladder region L with $a = 8$ and $b = 9$ is given. The small dots represent elements of L , the circles on the left and on the top of L represent the points $A^{(i)}$ and $E^{(i)}$, $i \in \{1, 2, 3\}$ that are specified by the minor $M = [1, 3, 4 \mid 1, 2, 4]$. The dotted lines indicate the lower boundary of $L^{(i)}$. Note that the point $(4, 9)$ is not an element of L . Therefore, in this example we have

$$d = n(a + b + 3) - \sum_{i=1}^n (u_i + v_i) - 1 = 44.$$



a. a ladder region with $a = 8$ and $b = 9$ b. a 10 dimensional face of $\Delta_{[1,3,4|1,2,4]}(\mathbf{Y})$

FIGURE 2.




FIGURE 3. Constructing a family of non-intersecting lattice paths, such that the i^{th} path stays above $L^{(i)}$, $i \in \{1, 2, 3\}$




FIGURE 4. The corresponding family of non-intersecting lattice paths, where the i^{th} path has north-east turns only in $L^{(i)}$ for $i \in \{1, 2, 3\}$

Proof. We will use results of Jürgen Herzog and Ngô Việt Trung. In Section 4 of [8], ladder determinantal rings are introduced and investigated.

We equip the indeterminates $x_{i,j}$, $i \in \{0, 1, \dots, b\}$ and $j \in \{0, 1, \dots, a\}$ with the following partial order:

$$x_{i,j} \leq x_{i',j'} \text{ if } i \geq i' \text{ and } j \leq j'.$$

A t -antichain in this partial order is a family of elements $x_{r_1,s_1}, x_{r_2,s_2}, \dots, x_{r_t,s_t}$ such that $r_1 < r_2 < \dots < r_t$ and $s_1 < s_2 < \dots < s_t$. Thus, a t -antichain corresponds to a sequence $(s_1, b - r_1), (s_2, b - r_2), \dots, (s_t, b - r_t)$ of t points in the ladder region associated with $\mathbf{Y}$, where each point lies strictly south-east of the previous ones.

Let D_t be the union of the last $u_t - 1$ rows and the last $v_t - 1$ columns of $\mathbf{Y}$. Let $\Delta_M(\mathbf{Y})$ be the simplicial complex whose k -dimensional faces are subsets of elements of $\mathbf{Y}$ of cardinality $k + 1$ which do not contain a t -antichain in D_t for $t \in \{1, 2, \dots, n + 1\}$. Let f_k be the number of k -dimensional faces of $\Delta_M(\mathbf{Y})$ for $k \geq 0$. Then, Corollary 4.3 of [8] states, that

$$\dim_K R_M(\mathbf{Y})_\ell = \sum_{k \geq 0} \binom{\ell - 1}{k} f_k.$$

In the following, we will find an expression for the numbers f_k involving certain families of non-intersecting lattice paths.

In Figure 2.b, a 10-dimensional face of $\Delta_{[1,3,4][1,2,4]}(\mathbf{Y})$ is shown, the elements of the face are indicated by bold dots. We will describe a modification of Viennot's ‘light and shadow procedure’ (with the sun in the top-left corner) that produces a family of n non-intersecting lattice paths such that the i^{th} path runs from $A^{(i)} = (0, u_{n+1-i})$ to $E^{(i)} = (a - v_{n+1-i}, b)$ and has north-east turns only in $L^{(i)}$, for $i \in \{1, 2, \dots, n\}$.

Imagine a sun in the top-left corner of the ladder region and a wall along the lower-right border $B^{(1)}$ of $L^{(1)}$. Then each lattice point (r, s) that is either in $B^{(1)}$ or corresponds to an element $x_{s,b-r}$ of the face casts a ‘shadow’ $\{(x, y) : x \geq r, y \leq s\}$.

The first path starts at $A^{(1)}$, goes along the north-east border of this shadow and terminates in $E^{(1)}$. In the left-most diagram of Figure 3, this is accomplished for the face shown in Figure 2.b.

In the next step, we remove the wall on $B^{(1)}$ and all the elements of the face which correspond to lattice points lying on the first path. Then the procedure is iterated. See Figure 3 for an example. Let P be the resulting family of non-intersecting lattice paths.

Now, for each $i \in \{1, 2, \dots, n\}$, we remove all elements of the face except those which correspond to north-east turns of the i^{th} path and do not lie on $B^{(i)}$. In the example, $(5, 8)$ is a north-east turn of the second path but lies on $B^{(2)}$, therefore the corresponding element $x_{1,5}$ of the face is removed. On the other hand, $(4, 5)$ lies on $B^{(1)}$, but is a north-east turn of the third path, so the corresponding element $x_{4,4}$ of the face is kept.

This set of north-east turns defines another family of non-intersecting lattice paths P' that has the property that the i^{th} path has north-east turns only in $L^{(i)}$ for $i \in \{1, 2, \dots, n\}$.

We now want to count the number of faces of $\Delta_M(\mathbf{Y})$ that reduce under ‘light and shadow’ to a given family of lattice paths P' with this property. Clearly, P' can be translated into a family P of non-intersecting lattice paths such that the i^{th}

path does not go below $B^{(i)}$ for $i \in \{1, 2, \dots, n\}$. Note that the number of lattice points on such a family P of paths is always equal to d , independently of the given face. Thus, if m is the number of north-east turns of P' , there are

$$\binom{d-m}{k+1-m}$$

families of non-intersecting lattice paths P that reduce to P' .

Hence, $f_k = \binom{d-m}{k+1-m} |\mathcal{T}_\ell^L(\mathbf{A} \mapsto \mathbf{E})|$ and we obtain

$$\begin{aligned} \sum_{\ell \geq 0} \dim_K R_M(\mathbf{Y})_\ell z^\ell &= \sum_{\ell \geq 0} \left(\sum_{k \geq 0} \binom{\ell-1}{k} f_k \right) z^\ell \\ &= \sum_{\ell \geq 0} \sum_{k \geq 0} \binom{\ell-1}{k} \left(\sum_{m=0}^{k+1} \binom{d-m}{k+1-m} |\mathcal{T}_m^L(\mathbf{A} \mapsto \mathbf{E})| \right) z^\ell \\ &= \sum_{m \geq 0} |\mathcal{T}_m^L(\mathbf{A} \mapsto \mathbf{E})| \sum_{\ell \geq 0} z^\ell \sum_{k \geq 0} \binom{\ell-1}{k} \binom{d-m}{d-k-1}, \end{aligned}$$

and if we sum the inner sum by means of the Vandermonde summation (see for example [7], Section 5.1, (5.27)),

$$\begin{aligned} \sum_{\ell \geq 0} \dim_K R_M(\mathbf{Y})_\ell z^\ell &= \sum_{m \geq 0} |\mathcal{T}_m^L(\mathbf{A} \mapsto \mathbf{E})| \sum_{\ell \geq 0} z^\ell \binom{d+\ell-m-1}{d-1} \\ &= \frac{\sum_{m \geq 0} |\mathcal{T}_m^L(\mathbf{A} \mapsto \mathbf{E})| z^m}{(1-z)^d}. \end{aligned}$$

□

4. LOG-CONCAVITY OF THE h -VECTOR IN THE CASE $M = [u_1 \mid v_1]$

In this paper we will settle Stanley's conjecture when R is a ladder determinantal ring cogenerated by M , where M is just a pair of integers, i.e., $n = 1$. We want to stress, however, that data strongly suggest that Conca and Herzog's conjecture is also true for arbitrary n .

By the preceding theorem, in the case we are going to tackle, the sum $\sum_{i=0}^s h_i x^i$ that appears in the conjecture is the generating function $\sum_{k \geq 0} |\mathcal{T}_k^L(A \mapsto E)| z^k$ of two-rowed arrays bounded by A and E which are in the ladder region L .

As the bounds A and E will not be of any significance throughout the rest of this paper, we will abbreviate $\mathcal{T}_k^L(A \mapsto E)$ to $\mathcal{T}_k^L$. We will show that the h -vector is log-concave by constructing an injection from $\mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$ into $\mathcal{T}_k^L \times \mathcal{T}_k^L$. This injection will involve some cut and paste operations that we now define:

Definition 4.1. Let A and X be two strictly increasing sequences of integers, such that the length of X is the length of A minus two, i.e., $A = (a_1, a_2, \dots, a_{k+1})$ and $X = (x_1, x_2, \dots, x_{k-1})$ for some $k \geq 1$. A *cutting point of A and X* is an index $l \in \{1, 2, \dots, k\}$ such that

$$\begin{aligned} (*) \quad & a_l < x_l, \\ & \text{and } x_{l-1} < a_{l+1}, \end{aligned}$$

where we require the inequalities to be satisfied only if all variables are defined. Hence, 1 is a cutting point if $a_1 < x_1$, and k is a cutting point if $x_{k-1} < a_{k+1}$.

The image of A and X obtained by cutting at l is

$$\begin{array}{cccccc|cccc} a_1 & a_2 & \dots & a_{l-1} & a_l & & x_l & x_{l+1} & \dots & x_{k-1} \\ \hline x_1 & x_2 & \dots & x_{l-1} & a_{l+1} & a_{l+2} & \dots & \dots & \dots & a_{k+1} \end{array}$$

Note that both the resulting sequences have length k .

Lemma 4.2. *Let $A = (a_1, a_2, \dots, a_{k+1})$ and $X = (x_1, x_2, \dots, x_{k-1})$ be strictly increasing sequences of integers, such that the length of X is the length of A minus two. Then there exists at least one cutting point of A and X .*

Proof. If $a_l \geq x_l$ for $l \in \{1, 2, \dots, k-1\}$ then $a_{k+1} > a_{k-1} \geq x_{k-1}$ and k is a cutting point. Otherwise, let l be minimal such that $a_l < x_l$. If $l = 1$ then 1 is a cutting point. Otherwise, because of the minimality of l , we have $a_{l+1} > a_{l-1} \geq x_{l-1}$, thus l is a cutting point. $\square$

Definition 4.3. Let $T = (T_1, T_2) \in \mathcal{T}_{k+1} \times \mathcal{T}_{k-1}$ be a pair of two-rowed arrays. Then a *top cutting point* of T is a cutting point of the top rows of T_1 and T_2 and a *bottom cutting point* of T is a cutting point of the bottom rows of T_1 and T_2 .

A pair (l, m) , where $l, m \in \{1, 2, \dots, k\}$, such that l is a top cutting point and m is a bottom cutting point of T_1 and T_2 is a *cutting point* of T . Cutting the top rows of T at l and the bottom rows at m we obtain the *image* of T . Note that both of the two-rowed arrays in the image have length k . More pictorially, if $l < m$,

$$\begin{array}{cccccc|cccc} a_1 & \dots & a_l & x_l & \dots & x_{m-1} & \dots & x_{k-1} \\ b_1 & \dots & b_{l+1} & \dots & b_m & y_m & \dots & y_{k-1} \\ \hline x_1 & \dots & x_{l-1} & a_{l+1} & \dots & a_m & \dots & a_{k+1} \\ y_1 & \dots & y_l & \dots & y_{m-1} & b_{m+1} & \dots & b_{k+1}, \end{array}$$

and similarly if $l \geq m$.

For $T = (T_1, T_2) \in \mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$, the pair (l, m) is an *allowed cutting point* of T , if both of the two-rowed arrays in the obtained image are in L .

In Lemma 5.1 we will prove that every pair of two-rowed arrays in $\mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$ has at least one allowed cutting point. This motivates the following definition:

Definition 4.4. Let $T = (T_1, T_2) \in \mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$ a pair of two-rowed arrays as before. Consider all allowed cutting points $(\bar{l}, \bar{m})$ of T . Select those with $|\bar{l} - \bar{m}|$ minimal. Among those, let (l, m) be the pair which comes first in the lexicographic order. Then we call (l, m) the *optimal cutting point* of T .

Now we are ready to state our main theorem, which implies that Stanley's conjecture is true, when R is a ladder determinantal ring cogenerated by a pair of integers M :

Theorem 4.5. *Let L be a ladder region. Let $T \in \mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$. Define $I(T)$ to be the pair of two-rowed arrays obtained by cutting T at its optimal cutting point. Then I is well-defined and an injection from $\mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$ into $\mathcal{T}_k^L \times \mathcal{T}_k^L$.*

Corollary 4.6. *The h -vector of the ladder determinantal ring cogenerated by $M = [u_1 \mid v_1]$ is log-concave.*

Proof of the corollary. By Theorem 3.1, the h -vector of this ring is equal to the generating function $\sum_{k \geq 0} |\mathcal{T}_k^L(A \mapsto E)| z^k$ of two-rowed arrays bounded by $A = (0, u_1 - 1)$ and $E = (a - v_1 + 1, b)$ which are in the ladder region L . By the

preceding theorem, there is an injection from $\mathcal{T}_{k+1}^L(A \mapsto E) \times \mathcal{T}_{k-1}^L(A \mapsto E)$ into $\mathcal{T}_k^L(A \mapsto E) \times \mathcal{T}_k^L(A \mapsto E)$, thus

$$|\mathcal{T}_{k+1}^L(A \mapsto E)| \cdot |\mathcal{T}_{k-1}^L(A \mapsto E)| \leq |\mathcal{T}_k^L(A \mapsto E)|^2.$$

□

We will split the proof of Theorem 4.5 in two parts. In Section 5 we show that the mapping I is well-defined, that is, for any pair of two-rowed arrays $T \in \mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$ there is an allowed cutting point. Finally, in Section 6, we show that I is indeed an injection.

5. THE MAPPING I IS WELL-DEFINED

Lemma 5.1. *Let L be a ladder region. Then for every pair of two-rowed arrays in $\mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$ there is an allowed cutting point (l, m) .*

For the proof of this lemma, we have to introduce some more notation: Let $(T_1, T_2) \in \mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$ with $T_1 = \begin{pmatrix} a_1 & a_2 & \dots & a_{k+1} \\ b_1 & b_2 & \dots & b_{k+1} \end{pmatrix}$ and $T_2 = \begin{pmatrix} x_1 & x_2 & \dots & x_{k-1} \\ y_1 & y_2 & \dots & y_{k-1} \end{pmatrix}$. We say that Inequality $(\overline{top})$ holds for an interval $[c, d]$ if

$$(\overline{top}) \quad \overline{L}(a_j) \geq y_{j-1},$$

for $j \in [c, d]$. Inequality $(\underline{top})$ holds for an interval $[c, d]$ if

$$(\underline{top}) \quad \underline{L}(a_j) \leq y_{j-1},$$

for $j \in [c, d]$. Similarly, Inequality $(\overline{bottom})$ holds for an interval $[c, d]$ if

$$(\overline{bottom}) \quad \overline{L}(x_{j-1}) \geq b_j,$$

for $j \in [c, d]$. Inequality $(\underline{bottom})$ holds for an interval $[c, d]$ if

$$(\underline{bottom}) \quad \underline{L}(x_{j-1}) \leq b_j,$$

for $j \in [c, d]$, where $\underline{L}$ and $\overline{L}$ are as in Definition 2.1. We say that any of these inequalities holds for a cutting point (l, m) if it holds for the interval $[l+1, m]$ if $l < m$ and for the interval $[m+1, l]$ if $m < l$. Clearly, a cutting point (l, m) is allowed if and only if all of these inequalities hold for it.

Most of the work is done by the following lemma:

Lemma 5.2. *Let $T = (T_1, T_2) \in \mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$, $T_1 = \begin{pmatrix} a_1 & a_2 & \dots & a_{k+1} \\ b_1 & b_2 & \dots & b_{k+1} \end{pmatrix}$ and $T_2 = \begin{pmatrix} x_1 & x_2 & \dots & x_{k-1} \\ y_1 & y_2 & \dots & y_{k-1} \end{pmatrix}$. Let $\underline{l}$ and $\overline{l}$ be top cutting points, such that there is no top cutting point in the closed interval $[\underline{l}+1, \overline{l}-1]$. Similarly, let $\underline{m}$ and $\overline{m}$ be bottom cutting points, such that there is no bottom cutting point in the closed interval $[\underline{m}+1, \overline{m}-1]$. Then for both of the intervals $[\underline{l}+1, \overline{l}]$ and $[\underline{m}+1, \overline{m}]$,*

- either $(\overline{top})$ or $(\overline{bottom})$ hold,
- either $(\underline{top})$ or $(\underline{bottom})$ hold,
- either $(\overline{top})$ or $(\underline{top})$ hold,
- either $(\overline{bottom})$ or $(\underline{bottom})$ hold.

Let $l_{min}, l_{max}, m_{min}$ and m_{max} be the minimal and maximal top and bottom cutting points. Then we have

- $(\overline{top})$ and $(\underline{bottom})$ hold for $[2, \max(l_{min}, m_{min})]$ and
- $(\underline{top})$ and $(\overline{bottom})$ hold for $[\min(l_{max}, m_{max}), k]$.

Proof. Suppose that $(\overline{top})$ does not hold for the interval $[\underline{l} + 1, \bar{l}]$. We claim that in that case there is an index $j \in [\underline{l} + 1, \bar{l} - 1]$ such that $a_j < x_j$: For, by hypothesis there is an index $i \in [\underline{l} + 1, \bar{l}]$ such that $\overline{L}(a_i) < y_{i-1}$. We have $\overline{L}(a_i) < y_{i-1} \leq \overline{L}(x_{i-1})$ and because $\overline{L}$ is a weakly increasing function, $a_i < x_{i-1}$. It follows that $a_{i-1} < a_i < x_{i-1} < x_i$. Thus, if $i = \bar{l}$ we choose $j = i - 1$, otherwise $j = i$.

The same statement is true if $(\underline{bottom})$ does not hold for the interval $[\underline{l} + 1, \bar{l}]$: In this case there must be an index $i \in [\underline{l} + 1, \bar{l}]$ such that $\underline{L}(x_{i-1}) > b_i$. We conclude that $\underline{L}(a_i) \leq b_i < \underline{L}(x_{i-1})$ and thus $a_i < x_{i-1}$.

Next, we will use induction to prove that

$$(**) \quad a_l < x_l$$

$$\text{and } a_{l+1} \leq x_{l-1}$$

for $l \in [\underline{l} + 1, \bar{l} - 1]$. We will first do an induction on l to establish the claim for $l \in [j, \bar{l} - 1]$.

We start the induction at $l = j$: Above we already found that $a_j < x_j$. Therefore we must have $a_{j+1} \leq x_{j-1}$, because otherwise j would satisfy $(*)$ and hence were a top cutting point.

Now suppose that $(**)$ holds for a particular $l < \bar{l} - 1$. Then $a_{l+1} \leq x_{l-1} < x_{l+1}$, and, because there is no top cutting point at $l + 1$, we have $a_{l+2} \leq x_l$.

Similarly, to establish $(**)$ for $l \in [\underline{l} + 1, j]$ we do a reverse induction on l . Suppose that $(**)$ holds for a particular $l > \underline{l} + 1$. Then $a_{l-1} < a_{l+1} \leq x_{l-1}$, and, because there is no top cutting point at $l - 1$, we have $a_l \leq x_{l-2}$.

Thus we obtain

$$\overline{L}(x_{\bar{l}-1}) \geq \overline{L}(x_{\bar{l}-2}) \geq \overline{L}(a_{\bar{l}}) \geq b_{\bar{l}}, \text{ and}$$

$$\overline{L}(x_{l-1}) \geq \overline{L}(a_{l+1}) \geq b_{l+1} \geq b_l,$$

which means that $(\overline{bottom})$ holds for the interval $[\underline{l} + 1, \bar{l}]$.

Furthermore,

$$\underline{L}(a_{l+1}) \leq \underline{L}(a_{l+2}) \leq \underline{L}(x_l) \leq y_l, \text{ and}$$

$$\underline{L}(a_l) \leq \underline{L}(x_{l-2}) \leq y_{l-2} \leq y_{l-1},$$

which means that $(\underline{top})$ holds for the interval $[\underline{l} + 1, \bar{l}]$.

Next we show that $(\overline{top})$ and $(\underline{bottom})$ hold for the interval $[2, l_{min}]$: Assume that either of these inequalities does not hold for the interval $[2, l_{min}]$ and that $[2, l_{min}]$ does not contain a top cutting point except l_{min} . Then the above reverse induction implies that $a_1 \leq a_3 < x_1$, which means that 1 is a top cutting point. Thus, $l_{min} = 1$ and the interval $[2, l_{min}]$ is empty.

The other assertions are shown in a completely analogous fashion. $\square$

We are now ready to establish Lemma 5.1:

Proof of Lemma 5.1. Let $T = (T_1, T_2) \in \mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$. By Lemma 4.2 there is at least one cutting point (l, m) of T . Let $l_{min}, l_{max}, m_{min}$ and m_{max} be the minimal and maximal top and bottom cutting points of T as before.

If there is an index j which is a top *and* a bottom cutting point of T , then – trivially – (j, j) is an allowed cutting point. Otherwise, we have to show that there is a cutting point (l, m) for which $(\overline{top})$, $(\underline{top})$, $(\overline{bottom})$, and $(\underline{bottom})$ hold. Suppose that this is not the case.

For the inductive proof which follows, we have to introduce a convenient indexing scheme for the sequence of top and bottom cutting points. Let

$$\begin{aligned} m_{1,0} &= \max\{m : m < l_{\min} \text{ and } m \text{ is a bottom cutting point}\}, \\ m_{i,0} &= \max\{m : m < l_{i-1,1} \text{ and } m \text{ is a bottom cutting point}\} \quad \text{for } i > 1, \\ \text{and } l_{i,0} &= \max\{l : l < m_{i,1} \text{ and } l \text{ is a top cutting point}\} \quad \text{for } i \geq 1, \end{aligned}$$

where $m_{i,j+1}$ is the bottom cutting point directly after $m_{i,j}$, and $l_{i,j+1}$ is the top cutting point directly after $l_{i,j}$. Furthermore, we set $l_{0,1} = l_{\min}$.

More pictorially, we have the following sequence of top and bottom cutting points for $i \geq 1$:

$$\cdots < m_{i,0} < l_{i-1,1} < l_{i-1,2} < \cdots < l_{i,0} < m_{i,1} < m_{i,2} < \cdots < m_{i+1,0} < \cdots$$

If $m_{\min} > l_{\min}$, then $m_{1,0}$ does not exist, of course. Note that there are no bottom cutting points between $l_{i,1}$ and $l_{i+1,0}$, and there are no top cutting points between $m_{i,1}$ and $m_{i+1,0}$.

Suppose first that $m_{\min} < l_{\min}$. By induction on i , we will show that $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ hold for the cutting points $(l_{i-1,1}, m_{i,0})$, where $i \geq 1$. By Lemma 5.2 we know that $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ are satisfied for the cutting point $(l_{\min}, m_{1,0})$, because $[m_{1,0} + 1, l_{\min}] \subseteq [2, l_{\min}]$. It remains to perform the induction step, which we will divide into five simple steps.

Step 1. $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ hold for the interval $[m_{i,0} + 1, l_{i-1,1}]$. This is just a restatement of the induction hypothesis, i.e., that $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ hold for the cutting point $(l_{i-1,1}, m_{i,0})$.

Step 2. Either $(\overline{\text{bottom}})$ or $(\overline{\text{top}})$ does not hold for the interval $[m_{i,0} + 1, m_{i,1}]$. Because of Step 1, not both of $(\overline{\text{bottom}})$ and $(\overline{\text{top}})$ can hold for $(l_{i-1,1}, m_{i,0})$, lest this was an allowed cutting point. Thus either $(\overline{\text{bottom}})$ or $(\overline{\text{top}})$ does not hold for $[m_{i,0} + 1, l_{i-1,0} + 1]$. This interval is contained in $[m_{i,0} + 1, m_{i,1}]$, thus the inequalities $(\overline{\text{bottom}})$ and $(\overline{\text{top}})$ cannot hold on this interval either.

Step 3. $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ hold for $[l_{i,0} + 1, m_{i,1}]$. Suppose that $(\overline{\text{bottom}})$ does not hold for $[m_{i,0} + 1, m_{i,1}]$. Then, by Lemma 5.2 we obtain that $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ hold for $[m_{i,0} + 1, m_{i,1}]$, because this interval contains no bottom cutting points except $m_{i,1}$. The same is true, if $(\overline{\text{top}})$ does not hold for $[m_{i,0} + 1, m_{i,1}]$. Because $[l_{i,0} + 1, m_{i,1}]$ is a subset of this interval, $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ hold for the cutting point $(l_{i,0}, m_{i,1})$, or, equivalently, for the interval $[l_{i,0} + 1, m_{i,1}]$.

Step 4. Either $(\overline{\text{bottom}})$ or $(\overline{\text{top}})$ does not hold for $[l_{i,0} + 1, l_{i,1}]$. Because of Step 3, not both of $(\overline{\text{bottom}})$ and $(\overline{\text{top}})$ can hold for the cutting point $(l_{i,0}, m_{i,1})$, nor for the greater interval $[l_{i,0} + 1, l_{i,1}]$.

Step 5. $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ hold for $[m_{i+1,0} + 1, l_{i,1}]$. The interval $[l_{i,0} + 1, l_{i,1}]$ does not contain a top cutting point except $l_{i,1}$, thus by Lemma 5.2 and Step 4 we see that $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ hold. Finally, because $[m_{i+1,0} + 1, l_{i,1}] \subset [l_{i,0} + 1, l_{i,1}]$, $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ hold for the cutting point $(l_{i,1}, m_{i+1,0})$.

If $l_{\max} > m_{\max}$, then we encounter a contradiction: Let r be such that $m_{r,0} = m_{\max}$. We have just shown that $(\overline{\text{top}})$ and $(\overline{\text{bottom}})$ hold for the cutting point $(l_{r-1,1}, m_{r,0})$. Furthermore, by Lemma 5.2, $(\overline{\text{bottom}})$ and $(\overline{\text{top}})$ hold for $[m_{r,0}, k]$

and thus also for $(l_{r-1,1}, m_{r,0})$. Hence, this would be an allowed cutting point, contradicting our hypothesis.

If $l_{max} < m_{max}$, let r be such that $l_{r,0} = l_{max}$. By the induction (Step 3) we find that $(\overline{top})$ and $(\overline{bottom})$ hold for the cutting point $(l_{r,0}, m_{r,1})$. Again, because of Lemma 5.2, we know that $(\overline{bottom})$ and $(\overline{top})$ holds for $[l_{r,0}, k]$ and thus also for $(l_{r,0}, m_{r,1})$. Hence, we had an allowed cutting point in this case also.

The case that $m_1 > l_1$ is completely analogous. $\square$

6. THE MAPPING I IS AN INJECTION

Lemma 6.1. *The mapping I defined above is an injection.*

Proof. Suppose that $I(T) = I(T')$ for $T = (T_1, T_2)$ and $T' = (T'_1, T'_2)$, such that T and T' are elements of $\mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$. Let (l, m) be the optimal cutting point of T , and let (l', m') be the optimal cutting point of T' .

Observe that we can assume $\min(l, m, l', m') = 1$, because the elements of T and T' with index less than or equal to this minimum retain their position in $I(T)$. Likewise, we can assume that $\max(l, m, l', m') = k$.

Furthermore, we can assume that $l \leq l'$, otherwise we exchange the meaning of T and T' . Thus, we have to consider the following twelve situations:

- (1) $1 = l \leq l' \leq m \leq m' = k$
- (2) $1 = l \leq l' \leq m' \leq m = k$
- (3) $1 = l \leq m \leq l' \leq m' = k$
- (4) $1 = l \leq m \leq m' \leq l' = k$
- (5) $1 = l \leq m' \leq l' \leq m = k$
- (6) $1 = l \leq m' \leq m \leq l' = k$
- (7) $1 = m \leq l \leq l' \leq m' = k$
- (8) $1 = m \leq l \leq m' \leq l' = k$
- (9) $1 = m \leq m' \leq l \leq l' = k$
- (10) $1 = m' \leq l \leq l' \leq m = k$
- (11) $1 = m' \leq l \leq m \leq l' = k$
- (12) $1 = m' \leq m \leq l \leq l' = k$

We shall divide these twelve cases into two portions according to whether $l \leq m$ or not.

A: $l \leq m$. In the Cases (1)–(6), (10) and (11) we have $l \leq m$, thus the pair of two-rowed arrays $T = (T_1, T_2) \in \mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$ looks like

$$\begin{array}{ccccccc} a_1 & \dots & a_l & | & a_{l+1} & \dots & a_{k+1} \\ b_1 & \dots & & & b_m & | & b_{m+1} \dots b_{k+1} \\ \hline x_1 & \dots & x_{l-1} & | & x_l & \dots & x_{k-1} \\ y_1 & \dots & & & y_{m-1} & | & y_m \dots y_{k-1} \end{array}$$

Cutting at (l, m) we obtain $I(T) \in \mathcal{T}_k^L \times \mathcal{T}_k^L$:

$$\begin{array}{ccccccc} a_1 & \dots & a_l & | & x_l & \dots & x_{k-1} \\ b_1 & \dots & & & b_m & | & y_m \dots y_{k-1} \\ \hline x_1 & \dots & x_{l-1} & | & a_{l+1} & \dots & a_{k+1} \\ y_1 & \dots & & & y_{m-1} & | & b_{m+1} \dots b_{k+1} \end{array}$$

If $l = 1$, then the top row of the second array in $I(T)$ is $(a_2, a_3, \dots, a_{k+1})$, if $m = k$, then the bottom row of the first array in $I(T)$ is $(b_1, b_2, \dots, b_k)$.

Case (1), $1 = l \leq l' \leq m \leq m' = k$. Given that $I(T) = I(T')$, the pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \begin{array}{cccccccccccccccc} a_1 & | & x_1 & \dots & x_{l'-1} & \vdots & a_{l'+1} & \dots & \dots & \dots & \dots & \dots & \dots & a_{k+1} \\ b_1 & \dots & \dots & \dots & \dots & \dots & \dots & \dots & b_m & | & y_m & \dots & y_{k-1} & \vdots & b_{k+1} \\ \hline a_2 & \dots & a_{l'} & \vdots & x_{l'} & \dots & \dots & \dots & \dots & \dots & x_{k-1} & \dots & \dots & \dots & \\ y_1 & \dots & \dots & \dots & \dots & \dots & y_{m-1} & | & b_{m+1} & \dots & b_k & \vdots & \dots & \dots & \end{array}$$

The vertical dots indicate the cut (l', m') which would result in $I(T')$. We show that the cutting point $(l, m) = (1, m)$, indicated above by the vertical lines, is in fact an allowed cutting point for T' : Cutting at $(1, m)$ yields

$$(\tilde{T}) \quad \begin{array}{cccccccccccccccc} a_1 & | & a_2 & \dots & a_{l'} & \vdots & x_{l'} & \dots & x_{m-1} & \dots & x_{k-1} & \dots & \dots & \dots & \\ b_1 & \dots & \dots & \dots & \dots & \dots & \dots & \dots & b_m & | & b_{m+1} & \dots & b_k & \vdots & \\ \hline x_1 & \dots & x_{l'-1} & \vdots & a_{l'+1} & \dots & a_m & \dots & \dots & \dots & a_{k+1} & \dots & \dots & \dots & \\ y_1 & \dots & \dots & \dots & \dots & \dots & y_{m-1} & | & y_m & \dots & y_{k-1} & \vdots & b_{k+1} & \dots & \end{array}$$

Note, that this is the same pair of two-rowed arrays we obtain by cutting T at (l', m') . We have to check that the pair of two-rowed arrays $(\tilde{T})$ is in the ladder region.

Clearly,

$$(a_2, b_2), (a_3, b_3), \dots, (a_{l'}, b_{l'}) \text{ and } (x_1, y_1), (a_2, b_2), \dots, (x_{l'-1}, y_{l'-1})$$

are in the ladder region, because these pairs appear also in T . Furthermore, the pairs

$$(x_{l'}, b_{l'+1}), (x_{l'+1}, b_{l'+2}), \dots, (x_{m-1}, b_m) \\ \text{and } (a_{l'+1}, y_{l'}), (a_{l'+2}, y_{l'+1}), \dots, (a_m, y_{m-1})$$

appear in $I(T)$ and are therefore in the ladder region, too. All the other pairs, i.e.,

$$(a_1, b_1) \text{ and } (x_m, b_{m+1}), (x_{m+1}, b_{m+2}), \dots, (x_{k-1}, b_k), \\ (a_{m+1}, y_m), (a_{m+2}, y_{m+1}), \dots, (a_k, y_{k-1}) \text{ and } (a_{k+1}, b_{k+1}),$$

are unaffected by the cut and appear in T' .

Thus we have that (l, m) and (l', m') are allowed cuts for T and T' . We required that (l, m) is optimal for T and that (l', m') is optimal for T' , therefore we must have $l = l'$ and $m = m'$.

In all the other cases the reasoning is very similar. Thus we only print the pairs of two-rowed arrays T' and $\tilde{T}$ and leave it to the reader to check that $\tilde{T}$ is in the ladder region.

Case (2), $1 = 1 \leq l' \leq m' \leq m = k$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \begin{array}{c} a_1 \mid x_1 \quad \dots \quad x_{l'-1} \vdots a_{l'+1} \quad \dots \quad a_{k+1} \\ b_1 \quad \dots \quad b_{m'} \vdots y_{m'} \quad \dots \quad y_{k-1} \mid b_{k+1} \\ \hline a_2 \quad \dots \quad a_{l'} \vdots x_{l'} \quad \dots \quad x_{k-1} \\ y_1 \quad \dots \quad y_{m'-1} \vdots b_{m'+1} \quad \dots \quad b_k \mid. \end{array}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \begin{array}{c} a_1 \mid a_2 \quad \dots \quad a_{l'} \vdots x_{l'} \quad \dots \quad x_{m'-1} \quad \dots \quad x_{k-1} \\ b_1 \quad \dots \quad b_{m'} \vdots y_{m'} \quad \dots \quad y_{k-1} \mid \\ \hline x_1 \quad \dots \quad x_{l'-1} \vdots a_{l'+1} \quad \dots \quad a_{m'} \quad \dots \quad a_{k+1} \\ y_1 \quad \dots \quad y_{m'-1} \vdots b_{m'+1} \quad \dots \quad b_k \mid b_{k+1}. \end{array}$$

Case (3), $1 = 1 \leq m \leq l' \leq m' = k$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \begin{array}{c} a_1 \mid x_1 \quad \dots \quad x_{l'-1} \vdots a_{l'+1} \quad \dots \quad a_{k+1} \\ b_1 \quad \dots \quad b_m \mid y_m \quad \dots \quad y_{k-1} \vdots b_{k+1} \\ \hline a_2 \quad \dots \quad a_{l'} \vdots x_{l'} \quad \dots \quad x_{k-1} \\ y_1 \quad \dots \quad y_{m-1} \mid b_{m+1} \quad \dots \quad b_k \vdots. \end{array}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \begin{array}{c} a_1 \mid a_2 \quad \dots \quad a_{l'} \vdots x_{l'} \quad \dots \quad x_{k-1} \\ b_1 \quad \dots \quad b_m \mid b_{m+1} \quad \dots \quad b_k \\ \hline x_1 \quad \dots \quad x_{l'-1} \vdots a_{l'+1} \quad \dots \quad a_{k+1} \\ y_1 \quad \dots \quad y_{m-1} \mid y_m \quad \dots \quad y_{k-1} \vdots b_{k+1}. \end{array}$$

Case (4), $1 = 1 \leq m \leq m' \leq l' = k$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \begin{array}{c} a_1 \mid x_1 \quad \dots \quad x_{k-1} \vdots a_{k+1} \\ b_1 \quad \dots \quad b_m \mid y_m \quad \dots \quad y_{m'-1} \vdots b_{m'+1} \quad \dots \quad b_{k+1} \\ \hline a_2 \quad \dots \quad a_k \vdots \\ y_1 \quad \dots \quad y_{m-1} \mid b_{m+1} \quad \dots \quad b_{m'} \vdots y_{m'} \quad \dots \quad y_{k-1}. \end{array}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \begin{array}{c} a_1 \mid a_2 \quad \dots \quad a_k \vdots \\ b_1 \quad \dots \quad b_m \mid b_{m+1} \quad \dots \quad b_{m'} \vdots y_{m'} \quad \dots \quad y_{k-1} \\ \hline x_1 \quad \dots \quad x_{k-1} \vdots a_{k+1} \\ y_1 \quad \dots \quad y_{m-1} \mid y_m \quad \dots \quad y_{m'-1} \vdots b_{m'+1} \quad \dots \quad b_{k+1}. \end{array}$$

Case (5), $1 = 1 \leq m' \leq l' \leq m = k$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \begin{array}{c} a_1 \mid x_1 \dots x_{l'-1} \vdots a_{l'+1} \dots a_{k+1} \\ b_1 \dots b_{m'} \vdots y_{m'} \dots y_{k-1} \mid b_{k+1} \\ \hline \begin{array}{c} a_2 \dots a_{l'} \vdots x_{l'} \dots x_{k-1} \\ y_1 \dots y_{m'-1} \vdots b_{m'+1} \dots b_k \mid \end{array} \end{array}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \begin{array}{c} a_1 \mid a_2 \dots a_{l'} \vdots x_{l'} \dots x_{k-1} \\ b_1 \dots b_{m'} \vdots y_{m'} \dots y_{k-1} \mid \\ \hline \begin{array}{c} x_1 \dots x_{l'-1} \vdots a_{l'+1} \dots a_{k+1} \\ y_1 \dots y_{m'-1} \vdots b_{m'+1} \dots b_k \mid b_{k+1} \end{array} \end{array}$$

Case (6), $1 = 1 \leq m' \leq m \leq l' = k$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \begin{array}{c} a_1 \mid x_1 \dots x_{k-1} \vdots a_{k+1} \\ b_1 \dots b_{m'} \vdots y_{m'} \dots y_{m-1} \mid b_{m+1} \dots b_{k+1} \\ \hline \begin{array}{c} a_2 \dots a_k \vdots \\ y_1 \dots y_{m'-1} \vdots b_{m'+1} \dots b_m \mid y_m \dots y_{k-1} \end{array} \end{array}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \begin{array}{c} a_1 \mid a_2 \dots a_k \vdots \\ b_1 \dots b_{m'} \vdots y_{m'} \dots y_{m-1} \mid y_m \dots y_{k-1} \\ \hline \begin{array}{c} x_1 \dots x_{k-1} \vdots a_{k+1} \\ y_1 \dots y_{m'-1} \vdots b_{m'+1} \dots b_m \mid b_{m+1} \dots b_{k+1} \end{array} \end{array}$$

Case (10), $1 = m' \leq 1 \leq l' \leq m = k$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \begin{array}{c} a_1 \dots a_l \mid x_l \dots x_{l'-1} \vdots a_{l'+1} \dots a_{k+1} \\ b_1 \vdots y_1 \dots y_{k-1} \mid b_{k+1} \\ \hline \begin{array}{c} x_1 \dots x_{l-1} \mid a_{l+1} \dots a_{l'} \vdots x_{l'} \dots x_{k-1} \\ \vdots b_2 \dots b_k \mid \end{array} \end{array}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \begin{array}{c} a_1 \dots a_l \mid a_{l+1} \dots a_{l'} \vdots x_{l'} \dots x_{k-1} \\ b_1 \vdots y_1 \dots y_{k-1} \mid \\ \hline \begin{array}{c} x_1 \dots x_{l-1} \mid x_l \dots x_{l'-1} \vdots a_{l'+1} \dots a_{k+1} \\ \vdots b_2 \dots b_k \mid b_{k+1} \end{array} \end{array}$$

Case (11), $1 = \mathbf{m}' \leq l \leq \mathbf{m} \leq l' = \mathbf{k}$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \begin{array}{cccccccccccc} a_1 & \dots & a_l & | & x_l & \dots & x_{k-1} & \vdots & a_{k+1} \\ b_1 & \vdots & y_1 & \dots & y_{m-1} & | & b_{m+1} & \dots & b_{k+1} \\ \hline x_1 & \dots & x_{l-1} & | & a_{l+1} & \dots & a_k & \vdots \\ \vdots & b_2 & \dots & b_m & | & y_m & \dots & y_{k-1}. \end{array}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \begin{array}{cccccccccccc} a_1 & \dots & a_l & | & a_{l+1} & \dots & a_k & \vdots \\ b_1 & \vdots & y_1 & \dots & y_{m-1} & | & y_m & \dots & y_{k-1} \\ \hline x_1 & \dots & x_{l-1} & | & x_l & \dots & x_{k-1} & \vdots & a_{k+1} \\ \vdots & b_2 & \dots & b_m & | & b_{m+1} & \dots & b_{k+1}. \end{array}$$

B: $\mathbf{m} \leq l$. In the Cases (7)–(9) and (12) we have $m \leq l$, thus the pair of two-rowed arrays $T = (T_1, T_2) \in \mathcal{T}_{k+1}^L \times \mathcal{T}_{k-1}^L$ looks like

$$\begin{array}{cccccccccccc} a_1 & \dots & a_l & | & a_{l+1} & \dots & a_{k+1} \\ b_1 & \dots & b_m & | & b_{m+1} & \dots & b_{k+1} \\ \hline x_1 & \dots & x_{l-1} & | & x_l & \dots & x_{k-1} \\ y_1 & \dots & y_{m-1} & | & y_m & \dots & y_{k-1}. \end{array}$$

Cutting at (l, m) we obtain $I(T) \in \mathcal{T}_k^L \times \mathcal{T}_k^L$:

$$\begin{array}{cccccccccccc} a_1 & \dots & a_l & | & x_l & \dots & x_{k-1} \\ b_1 & \dots & b_m & | & y_m & \dots & y_{k-1} \\ \hline x_1 & \dots & x_{l-1} & | & a_{l+1} & \dots & a_{k+1} \\ y_1 & \dots & y_{m-1} & | & b_{m+1} & \dots & b_{k+1}. \end{array}$$

Case (7), $1 = \mathbf{m} \leq l \leq l' \leq \mathbf{m}' = \mathbf{k}$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \begin{array}{cccccccccccc} a_1 & \dots & a_l & | & x_l & \dots & x_{l'-1} & \vdots & a_{l'+1} & \dots & a_{k+1} \\ b_1 & | & y_1 & \dots & y_{m-1} & | & a_{l+1} & \dots & a_{l'} & \vdots & x_{l'} & \dots & x_{k-1} & \vdots & b_{k+1} \\ \hline x_1 & \dots & x_{l-1} & | & a_{l+1} & \dots & a_{l'} & \vdots & x_{l'} & \dots & x_{k-1} \\ | & b_2 & \dots & b_m & | & y_m & \dots & y_{k-1} & \vdots & b_k & \vdots. \end{array}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \begin{array}{cccccccccccc} a_1 & \dots & a_l & | & a_{l+1} & \dots & a_{l'} & \vdots & x_{l'} & \dots & x_{k-1} \\ b_1 & | & b_2 & \dots & y_{m-1} & | & x_l & \dots & x_{l'-1} & \vdots & a_{l'+1} & \dots & a_{k+1} \\ \hline x_1 & \dots & x_{l-1} & | & x_l & \dots & x_{l'-1} & \vdots & a_{l'+1} & \dots & a_{k+1} \\ | & y_1 & \dots & y_{m-1} & | & b_{m+1} & \dots & b_{k-1} & \vdots & b_{k+1}. \end{array}$$

Case (8), $1 = \mathbf{m} \leq \mathbf{l} \leq \mathbf{m}' \leq \mathbf{l}' = \mathbf{k}$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \frac{\begin{array}{ccccccc} a_1 & \dots & a_l & | & x_l & \dots & x_{k-1} & \vdots & a_{k+1} \\ b_1 & | & y_1 & \dots & y_{m'-1} & \vdots & b_{m'+1} & \dots & b_{k+1} \end{array}}{\begin{array}{ccccccc} x_1 & \dots & x_{l-1} & | & a_{l+1} & \dots & a_k & \vdots \\ | & b_2 & \dots & b_{m'} & \vdots & y_{m'} & \dots & y_{k-1} \end{array}}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \frac{\begin{array}{ccccccc} a_1 & \dots & a_l & | & a_{l+1} & \dots & a_k & \vdots \\ b_1 & | & b_2 & \dots & b_{m'} & \vdots & y_{m'} & \dots & y_{k-1} \end{array}}{\begin{array}{ccccccc} x_1 & \dots & x_{l-1} & | & x_l & \dots & x_{k-1} & \vdots & a_{k+1} \\ | & y_1 & \dots & y_{m'-1} & \vdots & b_{m'+1} & \dots & b_{k+1} \end{array}}$$

Case (9), $1 = \mathbf{m} \leq \mathbf{m}' \leq \mathbf{l} \leq \mathbf{l}' = \mathbf{k}$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \frac{\begin{array}{ccccccc} a_1 & \dots & a_l & | & x_l & \dots & x_{k-1} & \vdots & a_{k+1} \\ b_1 & | & y_1 & \dots & y_{m'-1} & \vdots & b_{m'+1} & \dots & b_{k+1} \end{array}}{\begin{array}{ccccccc} x_1 & \dots & x_{l-1} & | & a_{l+1} & \dots & a_k & \vdots \\ | & b_2 & \dots & b_{m'} & \vdots & y_{m'} & \dots & y_{k-1} \end{array}}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \frac{\begin{array}{ccccccc} a_1 & \dots & a_l & | & a_{l+1} & \dots & a_k & \vdots \\ b_1 & | & b_2 & \dots & b_{m'} & \vdots & y_{m'} & \dots & y_{k-1} \end{array}}{\begin{array}{ccccccc} x_1 & \dots & x_{l-1} & | & x_l & \dots & x_{k-1} & \vdots & a_{k+1} \\ | & y_1 & \dots & y_{m'-1} & \vdots & b_{m'+1} & \dots & b_{k+1} \end{array}}$$

Case (12), $1 = \mathbf{m}' \leq \mathbf{m} \leq \mathbf{l} \leq \mathbf{l}' = \mathbf{k}$. The pair T' can be expressed in terms of the entries of T as follows:

$$(T') \quad \frac{\begin{array}{ccccccc} a_1 & \dots & a_l & | & x_l & \dots & x_{k-1} & \vdots & a_{k+1} \\ b_1 & \vdots & y_1 & \dots & y_{m-1} & | & b_{m+1} & \dots & b_{k+1} \end{array}}{\begin{array}{ccccccc} x_1 & \dots & x_{l-1} & | & a_{l+1} & \dots & a_k & \vdots \\ \vdots & b_2 & \dots & b_m & | & y_m & \dots & y_{k-1} \end{array}}$$

Cutting at (l, m) yields

$$(\tilde{T}) \quad \frac{\begin{array}{ccccccc} a_1 & \dots & a_l & | & a_{l+1} & \dots & a_k & \vdots \\ b_1 & \vdots & y_1 & \dots & y_{m-1} & | & y_m & \dots & y_{k-1} \end{array}}{\begin{array}{ccccccc} x_1 & \dots & x_{l-1} & | & x_l & \dots & x_{k-1} & \vdots & a_{k+1} \\ \vdots & b_2 & \dots & b_m & | & b_{m+1} & \dots & b_{k+1} \end{array}}$$

□

REFERENCES

- [1] Shreeram Abhyankar, *Enumerative combinatorics of Young tableaux*, Marcel Dekker, New York, 1988.
- [2] Shreeram Abhyankar and Devadatta M. Kulkarni, *On Hilbertian ideals*, Linear Algebra and its Applications **116** (1989), 53–79.
- [3] Francesco Brenti, *Log-concave and unimodal sequences in algebra, combinatorics, and geometry: an update*, Contemporary Mathematics **178** (1994), 71–89.
- [4] W. Bruns and Jürgen Herzog, *On the computation of a -invariants*, Manuscripta mathematica **77** (1992), 201–213.

- [5] A. Conca, *Ladder determinantal rings*, Journal of Pure and Applied Algebra **98** (1995), 119–134.
- [6] A. Conca and Jürgen Herzog, *On the Hilbert function of determinantal rings and their canonical module*, Proceedings of the American Mathematical Society **122** (1994), 677–681.
- [7] R. L. Graham, Donald E. Knuth, and Oren Patashnik, *Concrete mathematics*, Addison-Wesley, Reading, Massachusetts, 1989.
- [8] Jürgen Herzog and Ngô Viêt Trung, *Gröbner bases and multiplicity of determinantal and Pfaffian ideals*, Advances in Mathematics **96** (1992), no. 1, 1–37.
- [9] Christian Krattenthaler and Martin Rubey, *A determinantal formula for the Hilbert series of one-sided ladder determinantal rings*, preprint (2001).
- [10] Devadatta M. Kulkarni, *Hilbert polynomial of a certain ladder determinantal ideal*, Journal of Algebraic Combinatorics **2** (1993), 57–71.
- [11] ———, *Counting of paths and coefficients of Hilbert polynomial of a determinantal ideal*, Discrete Mathematics **154** (1996), 141–151.
- [12] G. Niesi and L. Robbiano, *Disproving Hibi’s conjecture with CoCoA or projective curves with bad Hilbert functions*, Computational Algebraic Geometry (Boston) (F. Eyssette and A. Galligo, eds.), Progress in Mathematics, no. 109, Birkhäuser, 1993, pp. 195–201.
- [13] Richard P. Stanley, *Log-concave and unimodal sequences in algebra, combinatorics, and geometry*, Annals of the New York Academy of Sciences **576** (1989), 500–535.